

AD-A279 169 V PAGE

Form Approved  
OMB No 0704-0188Public Report  
GPO: 1994-0-3  
Distribution  
Davis High

Four per response, including the time for reviewing instructions, searching existing data sources, gathering of information, and completion of the report. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Avenue, Washington, DC 20503.

|   |  |   |   |
|---|--|---|---|
| 1. AGENCY USE ONLY (Leave blank)  |  | 2. REPORT DATE<br>May 5, 1994                           | 3. REPORT TYPE AND DATES COVERED<br>Technical                                   |
| 4. TITLE AND SUBTITLE<br>Synthesis and Characterization of a Neutral Six-coordinate Tris-adduct of an Indium(III) Trihalide. X-ray Crystal Structure of $\text{InCl}_3(\text{THF})_3$ .   |  |   | 5. FUNDING NUMBERS<br>N00014-89-J1545<br>R&T Code 4135008                       |
| 6. AUTHOR(S)<br>R. L. Wells*, S. S. Kher, R. A Baldwin, and P. S. White   |  |   |   |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)<br>Department of Chemistry<br>Duke University<br>Durham, NC 27708-0346   |  |   | 8. PERFORMING ORGANIZATION REPORT NUMBER<br>Technical Report<br>No. DU/DC/TR-41 |
| 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)<br>Office of Naval Research<br>300 North Quincy Street<br>Arlington, VA 22217-5000  |  |   | 10. SPONSORING / MONITORING AGENCY REPORT NUMBER<br>94-14542                    |
| 11. SUPPLEMENTARY NOTES<br>Accepted for Publication in <i>Polyhedron</i>  |  |   |   |
| 12a. DISTRIBUTION / AVAILABILITY STATEMENT<br>Approved for Public Release<br>Distribution Unlimited   |  |   | 12b. DISTRIBUTION CODE  |
| 13. ABSTRACT (Maximum 200 words)<br><p>The reaction of <math>\text{InCl}_3</math> with excess THF afforded the adduct <math>\text{InCl}_3(\text{THF})_3</math> in a good yield (77.1% based on <math>\text{InCl}_3</math>), and the X-ray crystal structure shows it to be a hexa-coordinated tris-adduct of indium(III) chloride. The crystal structure consists of discrete monomeric units with no abnormally short intermolecular separations. The geometry about the indium atom approximates to octahedral with the three chlorine atoms (mean In-Cl = 2.422 Å) and the oxygen atoms of each THF molecule (mean In-O = 2.254 Å) residing in the meridional conformation. Crystals of <math>\text{InCl}_3(\text{THF})_3</math> belong to the monoclinic space group <math>P 2_1/c</math> with unit cell parameters <math>a = 8.1863(2)\text{Å}</math>, <math>b = 12.5141(2)\text{Å}</math>, <math>c = 16.815(4)\text{Å}</math>, <math>\beta = 93.84(4)^\circ</math>, <math>V = 1718.8(7)\text{Å}^3</math> for <math>Z = 4</math>. Full-matrix least-squares refinement based on 1603 reflections with <math>I &gt; 2.5\sigma(I)</math> converged at <math>R = 0.062</math> (<math>R_w = 0.084</math>). IR spectroscopic measurements gave absorption bands near 1027 and 857 <math>\text{cm}^{-1}</math>, which are consistent with coordinated THF molecules. Variable temperature NMR studies showed that the tris-adduct was stable to disproportionation in solution between 20-60 °C.</p> |  |   |   |
| 14. SUBJECT TERMS<br>Indium Compound, Six Coordinate<br>Synthesis, Crystal Structure  |  |   | 15. NUMBER OF PAGES<br>12   |
|   |  |   | 16. PRICE CODE  |
| 17. SECURITY CLASSIFICATION OF REPORT<br>Unclassified   | 18. SECURITY CLASSIFICATION OF THIS PAGE<br>Unclassified | 19. SECURITY CLASSIFICATION OF ABSTRACT<br>Unclassified | 20. LIMITATION OF ABSTRACT<br>Unlimited   |

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)  
Prescribed by ANSI Std. Z39-18  
298-102

94 5 13 110

OFFICE OF NAVAL RESEARCH

Grant N00014-89-J-1545

R&T Code 4135008

Technical Report No. DU/DC/TR-41

**Synthesis and Characterization of a  
Neutral Six-coordinate Tris-adduct of an Indium(III) Trihalide.  
X-ray Crystal Structure of  $\text{InCl}_3(\text{THF})_3$ .**

by

Richard L. Wells\*, Shreyas S. Kher, Ryan A. Baldwin and Peter S. White

Accepted for Publication in *Polyhedron*

Duke University  
Department of Chemistry,  
P. M. Gross Chemical Laboratory  
Box 90346  
Durham, NC 27708-0346

05 May 1994

|                    |                                     |
|--------------------|-------------------------------------|
| Accession For      |                                     |
| NTIS GRA&I         | <input checked="" type="checkbox"/> |
| DTIC TAB           | <input type="checkbox"/>            |
| Unannounced        | <input type="checkbox"/>            |
| Justification      |                                     |
| By                 |                                     |
| Distribution/      |                                     |
| Availability Codes |                                     |
| Dist               | Avail and/or<br>Special             |
| A-1                |                                     |

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

**SYNTHESIS AND CHARACTERIZATION OF A NEUTRAL  
SIX-COORDINATE TRIS-ADDUCT OF AN INDIUM(III)  
TRIHALIDE. X-RAY CRYSTAL STRUCTURE OF  $\text{InCl}_3(\text{THF})_3$ .**

**RICHARD L. WELLS\*, SHREYAS S. KHER, and RYAN A. BALDWIN**

Department of Chemistry, Paul M. Gross Chemical Laboratory, Duke University,  
Durham, North Carolina 27708, U.S.A.

**PETER S. WHITE**

Department of Chemistry, University of North Carolina at Chapel Hill,  
Chapel Hill, North Carolina 27514, U.S.A.

(Revised February 21, 1994, accepted ..... 1994)

**Abstract**—The reaction of  $\text{InCl}_3$  with excess THF afforded the adduct  $\text{InCl}_3(\text{THF})_3$  in a good yield (77.1% based on  $\text{InCl}_3$ ), and the X-ray crystal structure shows it to be a hexa-coordinated tris-adduct of indium(III) chloride. The crystal structure consists of discrete monomeric units with no abnormally short intermolecular separations. The geometry about the indium atom approximates to octahedral with the three chlorine atoms (mean  $\text{In-Cl} = 2.422 \text{ \AA}$ ) and the oxygen atoms of each THF molecule (mean  $\text{In-O} = 2.254 \text{ \AA}$ ) residing in the meridional conformation. Crystals of  $\text{InCl}_3(\text{THF})_3$  belong to the monoclinic space group  $P 2_1/c$  with unit cell parameters  $a = 8.1863(2) \text{ \AA}$ ,  $b = 12.5141(2) \text{ \AA}$ ,  $c = 16.815(4) \text{ \AA}$ ,  $\beta = 93.84(4)^\circ$ ,  $V = 1718.8(7) \text{ \AA}^3$  for  $Z = 4$ . Full-matrix least-squares refinement based on 1603 reflections with  $I > 2.5\sigma(I)$  converged at  $R = 0.062$  ( $R_w = 0.084$ ). IR spectroscopic measurements gave absorption bands near  $1027$  and  $857 \text{ cm}^{-1}$ , which are consistent with coordinated THF molecules. Variable temperature NMR studies showed that the tris-adduct was stable to disproportionation in solution between  $20$ – $60^\circ \text{C}$ .

---

Recent investigations in our laboratory<sup>1</sup> and that of Barron and co-workers<sup>2</sup> have shown that dehalosilylation is a facile means of forming the In-P bond. Thus, the latter obtained a product from the 1:1 mole ratio reaction of  $\text{InCl}_3$  and  $\text{P}(\text{SiMe}_3)_3$  at low temperature which was proposed to be  $[\text{Cl}_2\text{InP}(\text{SiMe}_3)_2]_x$ ; whereas, we found that the 2:1 mole reaction at room temperature afforded  $[\text{Cl}_2\text{In}_3\text{P}_3(\text{SiMe}_3)_2]_n$  and excess  $\text{InCl}_3$  as the penta-coordinated bis-adduct  $\text{InCl}_3(\text{THF})_2$  (I).<sup>3</sup> Subsequent to the serendipitous isolation of I, a direct synthesis was attempted. It was found that the reaction of  $\text{InCl}_3$  with excess THF afforded a six-coordinate compound  $\text{InCl}_3(\text{THF})_3$  II which is the tris-adduct. Compound II has been previously proposed by Fairbrother et al., but no solid-state structure was obtained.<sup>4</sup> They used colligative property measurements to postulate that the existence of II in the solid state would be of a lattice nature. We offer the solid-state structure and IR spectroscopic data which provide conclusive evidence that the three THF molecules present in the molecular structure are directly coordinated to the indium atom. In addition, to our knowledge, the crystal structure of II is one of the few examples of a neutral hexa-coordinated tris-adduct of an indium(III) trihalide with a meridional conformation (vide infra).

## EXPERIMENTAL

### Synthesis

All manipulations of air and moisture sensitive materials were performed in a Vacuum Atmospheres HE-493 Dri-Lab containing an argon atmosphere or by general Schlenk techniques. Toluene and THF were distilled from sodium benzophenone ketyl under dry dinitrogen.  $\text{InCl}_3$  was purchased from Strem Chemicals and was used as received without further purification.  $^1\text{H}$  and  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra were recorded on a Varian XL-300 spectrometer operating at 300 and 75.4 MHz, respectively.  $^1\text{H}$  and  $^{13}\text{C}\{^1\text{H}\}$  spectra were referenced to TMS by using the residual protons or carbons of deuterated benzene at  $\delta$  7.15 or 128 ppm. NMR samples were prepared in 5-mm tubes

under argon. IR spectroscopic data was collected on a BOMEM Michelson MB-100 Fourier Transform Infrared Spectrometer. A KBr pellet was prepared by adding 15 mg of **II** to 20 mg of KBr in an agate mortar and pestle with grinding to mix thoroughly. The sample transmittance was observed from 440-4000  $\text{cm}^{-1}$ . Melting points (uncorrected) were obtained on a Thomas-Hoover Uni-melt apparatus and the melting point capillaries were flame-sealed under argon. Elemental analyses were performed by E+R Microanalytical Laboratory, Inc., Corona, NY. X-ray crystallographic data were obtained on a Rigaku AFC6/S diffractometer utilizing graphite-monochromated Mo-K $\alpha$  radiation ( $\lambda = 0.71073\text{\AA}$ ) in the Single Crystal X-ray Structure Center at the University of North Carolina at Chapel Hill.

#### $\text{InCl}_3(\text{THF})_3$ (**II**)

Inside a dry-box, a one-necked 200  $\text{cm}^3$  round-bottomed flask equipped with a reflux condenser, Teflon valve, and a stir-bar was charged with  $\text{InCl}_3$  (0.659 g, 2.98 mmol) and 50  $\text{cm}^3$  of tetrahydrofuran. The mixture was removed from the dry-box and heated at reflux with stirring for 24 h under argon. The resulting clear solution was split into two equal portions in the dry-box. After diluting one portion with 2  $\text{cm}^3$  of toluene, both were cooled to  $-30^\circ\text{C}$ . After 3 days, colorless crystals of **II** (1.005 g total from both fractions, 77.1% yield based on  $\text{InCl}_3$ ; m.p. =  $59 - 62^\circ\text{C}$ ) formed in both portions. Found: C, 32.73; H, 5.40. Calc. for  $\text{C}_{12}\text{H}_{24}\text{Cl}_3\text{InO}_3$ : C, 32.94; H, 5.53; IR (KBr,  $\text{cm}^{-1}$ ) 3460 (vs), 1600 (vs), 1457 (w), 1027 (vs), 857 (vs);  $^1\text{H}$  NMR:  $\delta$  1.26 (m, 4H,  $\text{CH}_2$ ), 3.74 (m, 4H, O- $\text{CH}_2$ ).  $^{13}\text{C}\{^1\text{H}\}$  NMR:  $\delta$  25.16 (s,  $\text{CH}_2$ ), 69.20 (s, O- $\text{CH}_2$ ).

#### Variable temperature $^1\text{H}$ NMR

An internal standard of THF in degassed deuterated benzene was prepared by adding 0.02  $\text{cm}^3$  of THF to 0.06  $\text{cm}^3$   $\text{C}_6\text{D}_6$  in an NMR insert tube. A sample of **II** was prepared in an NMR tube by dissolving 16.9 mg of **II** in 0.6  $\text{cm}^3$  of  $\text{C}_6\text{D}_6$ . Individual

$^1\text{H}$  spectra of the internal standard [  $\delta$  1.48 (m, 4H,  $\text{CH}_2$ ), 3.52 (m, 4H,  $\text{O-CH}_2$ ) ] and of the internal standard inside the sample tube [  $\delta$  1.27 (m, 4H,  $\text{CH}_2$ ), 1.47 (m, 4H,  $\text{CH}_2$ ), 3.52 (m, 4H,  $\text{O-CH}_2$ ), 3.72 (m, 4H,  $\text{O-CH}_2$ ) ] were recorded at 19.5  $^\circ\text{C}$ . The internal standard was then removed and the sample tube was sealed with a rubber septum. The sample was heated in 5 $^\circ$  increments over a temperature range from 19.5-59.5  $^\circ\text{C}$ , and a  $^1\text{H}$  spectrum was recorded at each temperature [ 24.5  $^\circ\text{C}$  :  $\delta$  1.20 (m, 4H,  $\text{CH}_2$ ), 3.72 (m, 4H,  $\text{O-CH}_2$ ); 29.5  $^\circ\text{C}$  : 1.20 (m, 4H,  $\text{CH}_2$ ), 3.73 (m, 4H,  $\text{O-CH}_2$ ); 34.5  $^\circ\text{C}$  : 1.21 (m, 4H,  $\text{CH}_2$ ), 3.73 (m, 4H,  $\text{O-CH}_2$ ); 39.5  $^\circ\text{C}$  : 1.23 (m, 4H,  $\text{CH}_2$ ), 3.73 (m, 4H,  $\text{O-CH}_2$ ); 44.5  $^\circ\text{C}$  : 1.24 (m, 4H,  $\text{CH}_2$ ), 3.73 (m, 4H,  $\text{O-CH}_2$ ); 49.5  $^\circ\text{C}$  : 1.25 (m, 4H,  $\text{CH}_2$ ), 3.73 (m, 4H,  $\text{O-CH}_2$ ); 54.5  $^\circ\text{C}$  : 1.26 (m, 4H,  $\text{CH}_2$ ), 3.74 (m, 4H,  $\text{O-CH}_2$ ); 59.5  $^\circ\text{C}$  : 1.27 (m, 4H,  $\text{CH}_2$ ), 3.74 (m, 4H,  $\text{O-CH}_2$ ) ].

#### X-ray structural solution and refinement

Crystallographic data are summarized in Table I. The crystal used was a colorless block which was mounted on a glass fiber with a viscous oil under a stream of cold dinitrogen. X-ray intensity data were recorded at -170  $^\circ\text{C}$ , and the structure was solved by direct methods. Full-matrix least-squares refinement with weights based upon counter statistics was performed. Hydrogen atoms were incorporated at their calculated positions using a riders model in the later iterations of refinement which converged at  $R = 0.062$  ( $R_w = 0.084$ ). A final difference-Fourier synthesis revealed no unusual features (max. 1.28, min. -1.34  $\text{e } \text{\AA}^{-3}$ ). Crystallographic calculations were performed using the NRCVAX<sup>5</sup> suite of structure determination programs. For all structure-factor calculations, neutral atom scattering factors and their anomalous dispersion corrections were taken from ref. 6. Interatomic distances and angles are given in Table II. An ORTEP<sup>7</sup> diagram showing the solid-state conformation and atom numbering scheme of **II** is presented in Fig. 1. Full information concerning conditions for crystallographic data collection and structure refinement, atomic coordinates, thermal and positional

parameters, and observed and calculated structure factors has been deposited with the Cambridge Crystallographic Data Centre.

## RESULTS AND DISCUSSION

Although there have been numerous examples of six-coordinate indium(III) trihalide compounds reported to date<sup>8-12</sup>, an extensive review of the literature finds limited X-ray crystal analyses of such systems.<sup>9-12</sup> The principal methods for compound identification and structure elucidation have been Infrared and Raman spectroscopy, as well as colligative property measurements. Tris-adducts of indium(III) trihalides which have been crystallographically characterized are:  $[\text{InX}_3\text{L}_3 \cdot \text{Y}_3]$ ,  $\text{X} = \text{Cl}$ ,  $\text{Y} = \text{dioxane}$ ,  $\text{L} = \text{H}_2\text{O}$  (III)<sup>9</sup>;  $\text{InX}_3\text{L}_3$ ,  $\text{X} = \text{Cl}$ ,  $\text{L} = \text{terpy}$  (2, 2', 2''-terpyridyl) (IV)<sup>10</sup>,  $\text{Me}_3\text{PO}$  (V)<sup>11</sup>,  $\text{Me}_2\text{SO}$  (VI)<sup>11</sup>,  $\text{dmf}$  (dimethylformamide) (VII)<sup>12</sup>,  $\text{dma}$  (dimethylacetamide) (VIII)<sup>12</sup>;  $\text{X} = \text{Br}$ ,  $\text{L} = \text{Me}_2\text{SO}$  (IX)<sup>11</sup>]. It is interesting to note that five of these seven (V - IX) were reported within the last five years.

The crystal structure of II consists of discrete monomeric units with the chlorine atoms and the THF oxygen atoms approximately octahedrally coordinated in a meridional conformation about the indium atom. Octahedral geometry is common to all of the above  $\text{InX}_3 \cdot \text{L}_3$  adducts, however, III and VI - VIII are in the facial conformation. The  $\text{Cl}(1) - \text{In} - \text{O}(21)$  angle at  $179.2(3)^\circ$  in II deviates only slightly from an exactly linear value and is considerably greater than the mean  $\text{Cl} - \text{In} - \text{Cl}$  angle ( $166.7^\circ$ ) in III. The  $\text{O}(11) - \text{In} - \text{O}(31)$  and the  $\text{Cl}(3) - \text{In} - \text{Cl}(2)$  angles at  $168.8(3)^\circ$  and  $169.5(1)^\circ$ , respectively, evidence the degree of distortion from perfect octahedral symmetry. The mean  $\text{In} - \text{Cl}$  bond length at  $2.421 \text{ \AA}$  in II is nearly identical to that of  $2.422 \text{ \AA}$  in III and shorter than the corresponding lengths of  $2.456 \text{ \AA}$  in IV and  $2.435\text{-}2.608 \text{ \AA}$  in V - VIII. The  $\text{In} - \text{O}(21)$  bond in II at  $2.304 \text{ \AA}$ , which is trans to  $\text{Cl}(1)$ , is considerably longer than the  $\text{In} - \text{O}(11)$  [ $2.235(1) \text{ \AA}$ ] and  $\text{In} - \text{O}(31)$  [ $2.223(1) \text{ \AA}$ ] bonds. This bond lengthening may be attributed to steric interactions between the THF molecules or a trans-ligand

influence where the donor ability of the Cl atom serves to weaken the In - O bond. The In - O bond lengthening is not observed, however, in V where the In - O bond lengths differ by only 0.007 Å. The two strong IR absorptions at 1027 and 857  $\text{cm}^{-1}$  correspond to the symmetric and asymmetric C-O-C stretching vibrations of the THF molecules and are near the values of 1011 and 850  $\text{cm}^{-1}$  characteristic for coordinated THF molecules.<sup>13</sup> Fairbrother et al. used vapor pressure measurements on homogeneous unsaturated solutions of  $\text{InCl}_3$  in THF to conclude that the bis-adduct I would predominate in solution, and that the tris-adduct II most likely exists only in the solid-state as a lattice compound. However, we concluded from variable temperature NMR studies that the tris-adduct is stable to disproportionation in deuterated benzene solution at temperatures ranging from 20-60 °C.

**Acknowledgement-** We wish to thank the Office of Naval Research and the AT&T Bell Laboratories Cooperative Research Fellowship Program for their financial support . We also wish to recognize Dr. Leonidas J. Jones, III for his contributions to the NMR studies.

## References

- 1) For example, see the following and references therein: (a) R. L. Wells, A. T. McPhail and M. F. Self, *Organometallics* 1992, **11**, 221. (b) R. L. Wells, A. T. McPhail, L. J. Jones and M. F. Self, *Polyhedron* 1993, **12**, 141.
- 2) M. D. Healy, P. E. Laibinis, P. D. Stupik and A. R. Barron, *J. Chem. Soc., Chem. Commun.* 1989, 359.
- 3) M. F. Self, A. T. McPhail and R. L. Wells, *Polyhedron* 1993, **12**, 455.
- 4) F. Fairbrother, N. Flitcroft and H. Prophet, *J. Less-Common Met.* 1960, **2**, 49.
- 5) E. J. Gabe, Y. L. Page, J. P. Charland, F. L. Lee and P. S. White, *J. Appl. Cryst.* 1989, **22**, 384.
- 6) *International Tables for X-ray Crystallography*, Vol. IV. Kynoch Press, Birmingham (1974).
- 7) C. K. Johnson, ORTEP - A Fortran Thermal Ellipsoid Plot Program, Technical Report ORNL-5138, Oak Ridge 1976.
- 8) For example, see the following and references therein: (a) A. J. Carty and D. G. Tuck, *J. Chem. Soc. (A), Dalton Trans.* 1966, 1081. (b) D. H. Brown and D. T. Stewart, *J. Inorg. Nucl. Chem.* **32**, 3751. (c) H. L. Chung and D. G. Tuck, *Can. J. Chem.* 1975, **53**, 3492.
- 9) S. H. Whitlow and E. J. Gabe, *Acta Cryst.* 1975, **B31**, 2534.
- 10) G. Beran, K. Dymock, H. A. Patel, A. J. Carty and P. M. Boorman, *Inorganic Chemistry* 1972, **11**, 896.
- 11) W. T. Robinson, C. J. Wilkins and Z. Zeying, *J. Chem. Soc., Dalton Trans.* 1988, 2187.
- 12) S. Jin, V. McKee, M. Nieuwenhuyzen, W. T. Robinson and C. J. Wilkins, *J. Chem. Soc., Dalton Trans.* 1993, 3111.
- 13) R. J. H. Clark, J. Lewis, D. J. Machin, R. S. Nyholm, *J. Chem. Soc.* 1963, 379.

Caption to Figure

Fig. 1. ORTEP diagram ( 30% probability ellipsoids ) showing the solid-state conformation and atom numbering scheme of  $\text{InCl}_3(\text{THF})_3$  (II). The hydrogen atoms are represented by small circles.




Fig. 1

Table I. Crystallographic Data and Measurements for  $\text{InCl}_3(\text{THF})_3$  (II)

| II                                      |   |
|---|---|
| molecular formula                       | $\text{C}_{12}\text{H}_{24}\text{Cl}_3\text{InO}_3$ |
| formula weight                          | 437.49  |
| crystal system                          | monoclinic  |
| space group                             | $P 2_1/c$   |
| a, Å                                    | 8.1863(2)   |
| b, Å                                    | 12.5141(2)  |
| c, Å                                    | 16.815(4)   |
| $\beta$ , deg                           | 93.84(4)  |
| V, Å <sup>3</sup>                       | 1718.8(7)   |
| Z                                       | 4   |
| radiation (wavelength, Å)               | Mo-K $\alpha$ (0.71073)                             |
| $\mu$ , cm <sup>-1</sup>                | 18.3  |
| temp, °C                                | -170  |
| D <sub>calcd</sub> , g cm <sup>-3</sup> | 1.691   |
| crystal dims., mm                       | 0.30 x 0.30 x 0.30                                  |
| T <sub>max</sub> ; T <sub>min</sub>     | 1.00:0.37   |
| scan type                               | $\omega$  |
| scan width, deg                         | 1.00  |
| $\Theta_{\text{max}}$ , deg             | 45  |
| no. of rflns recorded                   | 3401  |
| no. of non-equiv.                       | 2251  |
| rflns recorded                          |   |
| R <sub>merg</sub> (on I)                | 0.063   |
| no. of rflns retained,                  | 1603  |
| $I > 2.5\sigma(I)$                      |   |
| no. of params. refined                  | 173   |
| R; R <sub>w</sub> <sup>a</sup>          | 0.062; 0.084  |

Table I (continued)

|   | II            |
|---|---------------|
| goodness-of-fit <sup>b</sup>                    | 2.27          |
| max shift / esd. in final                       | 0.002         |
| least-squares cycle                             |               |
| final max, min $\Delta\rho$ , e/Å <sup>-3</sup> | 1.280; -1.340 |

<sup>a</sup> $R = \Sigma(|F_O| - |F_C|)/\Sigma|F_O|$ ;  $R_w = [\Sigma w (|F_O| - |F_C|)^2 / \Sigma w |F_O|^2]^{1/2}$ .

<sup>b</sup>Goodness-of-fit =  $[\Sigma w \Delta^2 / (N_{\text{observations}} - N_{\text{parameters}})]^{1/2}$ .

Table II. Interatomic Distances (Å) and Angles (°) for  $\text{InCl}_3(\text{THF})_3$  (II), with Estimated Standard Deviations in Parentheses

|                   |           |                   |          |
|-------------------|-----------|-------------------|----------|
| In(1)-Cl(1)       | 2.423(4)  | O(21)-C(22)       | 1.412(2) |
| In(1)-Cl(2)       | 2.422(4)  | O(21)-C(25)       | 1.500(2) |
| In(1)-Cl(3)       | 2.420(4)  | C(22)-C(23)       | 1.546(2) |
| In(1)-O(11)       | 2.235(1)  | C(23)-C(24)       | 1.56(3)  |
| In(1)-O(21)       | 2.304(9)  | C(24)-C(25)       | 1.487(2) |
| In(1)-O(31)       | 2.223(1)  | O(31)-C(32)       | 1.454(2) |
| O(11)-C(12)       | 1.497(2)  | O(31)-C(35)       | 1.494(2) |
| O(11)-C(15)       | 1.446(2)  | C(32)-C(33)       | 1.436(2) |
| C(12)-C(13)       | 1.50(3)   | C(33)-C(34)       | 1.566(2) |
| C(13)-C(14)       | 1.536(2)  | C(34)-C(35)       | 1.480(2) |
| C(14)-C(15)       | 1.503(2)  |                   |          |
| Cl(1)-In(1)-Cl(2) | 94.65(1)  | O(11)-C(12)-C(13) | 104.2(1) |
| Cl(1)-In(1)-Cl(3) | 95.57(1)  | C(12)-C(13)-C(14) | 103.0(1) |
| Cl(1)-In(1)-O(11) | 98.1(3)   | C(13)-C(14)-C(15) | 101.8(1) |
| Cl(1)-In(1)-O(21) | 179.2(3)  | O(11)-C(15)-C(14) | 108.5(1) |
| Cl(1)-In(1)-O(31) | 93.1(3)   | In(1)-O(21)-C(22) | 128.5(9) |
| Cl(2)-In(1)-Cl(3) | 169.52(1) | In(1)-O(21)-C(25) | 122.4(7) |
| Cl(2)-In(1)-O(11) | 91.69(2)  | C(22)-O(21)-C(25) | 108.6(1) |
| Cl(2)-In(1)-O(21) | 84.99(2)  | O(21)-C(22)-C(23) | 108.4(1) |
| Cl(2)-In(1)-O(31) | 87.7(3)   | C(22)-C(23)-C(24) | 99.2(1)  |
| Cl(3)-In(1)-O(11) | 89.2(3)   | C(23)-C(24)-C(25) | 103.4(2) |
| Cl(3)-In(1)-O(21) | 84.77(2)  | O(21)-C(25)-C(24) | 105.1(i) |
| Cl(3)-In(1)-O(31) | 82.6(3)   | In(1)-O(31)-C(35) | 127.6(9) |
| O(11)-In(1)-O(31) | 168.8(3)  | C(32)-O(31)-C(35) | 108.8(1) |
| O(21)-In(1)-O(31) | 86.2(4)   | O(31)-C(32)-C(33) | 103.6(1) |
| In(1)-O(11)-C(12) | 124.8(8)  | C(32)-C(33)-C(34) | 103.8(1) |
| In(1)-O(11)-C(15) | 127.5(8)  | C(33)-C(34)-C(35) | 102.0(1) |
| C(12)-O(11)-C(15) | 107.7(1)  | O(31)-C(35)-C(34) | 105.6(1) |